arXiv:1506.02487v1 [math.CA] 24 May 2015

Some approximation properties of bivariate Bleimann-Butzer and Hahn operators based on (p, q)-integers

M. Mursaleen and **Md. Nasiruzzaman**
Department of Mathematics, Aligarh Muslim University, Aligarh–202002, India
mursaleenm@gmail.com; nasir3489@gmail.com

Abstract

Recently, Mursaleen et al applied (p, q)-calculus in approximation theory and introduced (p, q)-analogue of Bernstein operators in [16]. In this paper, we construct and introduce a generalization of the bivariate Bleimann-Butzer-Hahn operators based on (p, q)-integers and obtain Korovkin type approximation theorem of these operators. Furthermore, we compute the rate of convergence of the operators by using the modulus of continuity and Lipschitz type maximal functions.

Keywords and phrases: (p, q)-integers; (p, q)-Bernstein operator; (p, q)-Bleimann-Butzer-Hahn operators; q-Bivariate Bleimann-Butzer-Hahn operators; Korovkin theorem; modulus of continuity; Lipschitz type maximal function.

AMS Subject Classification (2010): 41A10, 41A25, 41A36, 40A30.

1. Introduction and Preliminaries

In 1980 Bleimann, Butzer and Hahn (BBH) [6] introduced the following positive linear operators on the space of real functions defined on infinite interval $[0, \infty)$:

$$L_n(f;x) = \frac{1}{(1+x)^n} \sum_{k=0}^{n} f\left(\frac{k}{n-k+1}\right) \left[\begin{array}{c} n \\ k \end{array}\right] x^k, x \geq 0 \tag{1.1}$$

The approximation properties of the Bleimann-Butzer and Hahn operators were studied by many authors, e.g. [1, 2, 3, 6]. In approximation theory, q-type generalization of Bernstein polynomials was introduced and studied by Lupaş [13] and Phillips [21].

In [8],[5], the BBH-type operators based on q-integers are defined as follows:

$$L_n^q(f;x) = \frac{1}{\ell_n(x)} \sum_{k=0}^{n} f\left(\frac{[k]_q}{[n-k+1]_q q^k}\right) q^{\frac{k(k-1)}{2}} \left[\begin{array}{c} n \\ k \end{array}\right]_q x^k \tag{1.2}$$

where $\ell_n(x) = \prod_{s=0}^{n-1}(1+q^s x)$.

The bivariate case was introduced by D.D. Stancu [25] who studied the bivariate Bernstein polynomials and estimated the order of approximation for operators (1.2).

Very recently, Mursaleen et al applied (p, q)-calculus in approximation theory and introduced first (p, q)-analogue of Bernstein operators [16] and studied approximation properties of (p, q)-analogue of Bernstein-Stancu operators in [17]. They also have introduced, Bleimann-Butzer and Hahn operators [20] based on (p, q)-integers as follows:

$$L_n^{(p,q)}(f;x) = \frac{1}{\ell_n^{(p,q)}(x)} \sum_{k=0}^{n} f\left(\frac{p^{n-k+1}[k]_{p,q}}{[n-k+1]_{p,q} q^k}\right) p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} x^k \tag{1.3}$$

where, $x \geq 0,\ 0 < q < p \leq 1$

$$\frac{1}{\ell_n^{(p,q)}(x)} = \prod_{s=0}^{n-1}(p^s + q^s x), \tag{1.4}$$

and f is defined on semiaxis $\mathbb{R}_+$.
The Euler identity based on (p,q)-analogue is defined by

$$\prod_{s=0}^{n-1}(p^s + q^s x) = \sum_{k=0}^{n} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} x^k \tag{1.5}$$

Let us recall certain notations on (p,q)-calculus.
The (p,q) integers $[n]_{p,q}$ are defined by

$$[n]_{p,q} = \frac{p^n - q^n}{p - q},\ n = 0, 1, 2, \cdots,\ 0 < q < p \leq 1.$$

whereas q-integers are given by

$$[n]_q = \frac{1 - q^n}{1 - q},\ n = 0, 1, 2, \cdots,\ 0 < q < 1.$$

It is very clear that q-integers and (p,q)-integers are different, that is we cannot obtain (p,q) integers just by replacing q by $\frac{q}{p}$ in the definition of q-integers but if we put $p = 1$ in definition of (p,q) integers then q-integers become a particular case of (p,q) integers. Thus we can say that (p,q)-calculus can be taken as a generalization of q-calculus.

Now by some simple calculation and induction on n, we have (p,q)-binomial expansion as follows

$$\begin{aligned}(ax + by)_{p,q}^n &:= \sum_{k=0}^{n} p^{\frac{(n-k)(n-k-1)}{2}} q^{\frac{k(k-1)}{2}} \left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} a^{n-k} b^k x^{n-k} y^k \\ (x + y)_{p,q}^n &= (x + y)(px + qy)(p^2x + q^2y)\cdots(p^{n-1}x + q^{n-1}y) \\ (1 - x)_{p,q}^n &= (1 - x)(p - qx)(p^2 - q^2x)\cdots(p^{n-1} - q^{n-1}x),\end{aligned}$$

and the (p,q)-binomial coefficients are defined by

$$\left[\begin{array}{c} n \\ k \end{array}\right]_{p,q} = \frac{[n]_{p,q}!}{[k]_{p,q}![n-k]_{p,q}!}.$$

Again it can be easily verified that (p,q)-binomial expansion is different from q-binomial expansion and is not a replacement of q by $\frac{q}{p}$.

By some simple calculation, we have the following relation

$$q^k[n - k + 1]_{p,q} = [n + 1]_{p,q} - p^{n-k+1}[k]_{p,q}. \tag{1.6}$$

For details on q-calculus and (p,q)-calculus, one can refer [5, 26], [22, 23], respectively.

Now based on (p,q)-integers, we construct (p,q)-analogue of bivariate BBH-operators, and we call it as (p,q)-bivariate Bleimann-Butzer-Hahn operators and investigate its Korovokin type approximation properties, by using the following test functions

$$e_{ij} = \left(\frac{u}{1+u}\right)^i \left(\frac{v}{1+v}\right)^j \quad i, j = 0, 1, 2. \tag{1.7}$$

Also for a space of generalized Lipschitz-type maximal functions, we give a pointwise estimation.

2. Construction of operators

Let $\mathbb{R}_+^2=[0,\infty)\times[0,\infty)$, $f:\mathbb{R}_+^2\to\mathbb{R}$ and $0<q_{n_1},q_{n_2}<p_{n_1},p_{n_2}\leq 1$. We define the bivariate extension of the (p,q)-Bleimann-Butzer and Hahn-type operators based on (p,q)-integers as follows:

$$L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y)=\tfrac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\tfrac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\textstyle\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2} f\left(\tfrac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{[n_1-k_1+1]_{p_{n_1},q_{n_1}}q_{n_1}^{k_1}},\tfrac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right)$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c}n_1\\k_1\end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c}n_2\\k_2\end{array}\right]_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2} \qquad (2.1)$$

where

$$\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}=\prod_{s_1=0}^{n_1-1}(p_{n_1}^{s_1}+q_{n_1}^{s_1}x),\quad \frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}=\prod_{s_2=0}^{n_2-1}(p_{n_2}^{s_2}+q_{n_2}^{s_2}y).$$

It is easy to check that (2.1) is linear and positive. For $p_{n_1}=p_{n_2}=1$, the operators (2.1) are reduced to q-bivariate BBH operators [8, 9].

Lemma 2.1. *Let $e_{ij}:\mathbb{R}_+^2\to[0,1)$ be the two dimensional test functions defined in (1.7). Then the (p,q)-bivariate BBH operators defined in (2.1) satisfy the following identities.*

(i) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)=1$

(ii) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{10};x,y)=\frac{[n_1]_{P_{n_1},q_{n_1}}}{[n_2+1]_{p_{n_1},q_{n_1}}}\frac{x}{1+x}$

(iii) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{01};x,y)=\frac{[n_2]_{P_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}\frac{y}{1+y}$

(iv) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{20};x,y)=\frac{p_{n_1}q_{n_1}^2[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2}\frac{x^2}{(1+x)(p_{n_1}+q_{n_1}x)}+\frac{p_{n_1}^{n_1+1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2}\left(\frac{x}{1+x}\right)$

(v) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{02};x,y)=\frac{p_{n_2}q_{n_2}^2[n_2]_{p_{n_2},q_{n_2}}[n_2-1]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}^2}\frac{y^2}{(1+y)(p_{n_2}+q_{n_2}y)}+\frac{p_{n_2}^{n_2+1}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}^2}\left(\frac{y}{1+y}\right).$

Proof. (i) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)=\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_1=1}^{n_1}\sum_{k_2=1}^{n_2}$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_{n_1}}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c}n_1\\k_1\end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c}n_2\\k_2\end{array}\right]_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}$$

For $0<q<p\leq 1$, we have

$$\sum_{k=0}^{n}p^{\frac{(n-k)(n-k-1)}{2}}q^{\frac{k(k-1)}{2}}\left[\begin{array}{c}n\\k\end{array}\right]_{p,q}x^k=\prod_{k=0}^{n-1}(p^k+q^kx)=\frac{1}{\ell_n^{p,q}(x)}.$$

(ii) Let $u=\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{[n_1-k_1+1]_{p_{n_1},q_{n_1}}q_{n_1}^{k_1}}$. Then $\frac{u}{1+u}=\frac{[k_1]_{p_{n_1},q_{n_1}}p_{n_1}^{n_1-k_1+1}}{[n_1+1]_{p_{n_1},q_{n_1}}}$

$$L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{10};x,y)=\tfrac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\tfrac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\textstyle\sum_{k_1=1}^{n_1}\sum_{k_2=1}^{n_2}\tfrac{[k_1]_{p_{n_1},q_{n_1}}p_{n_1}^{n_1-k_1+1}}{[n_1+1]_{p_{n_1},q_{n_1}}}\tfrac{[k_2]_{p_{n_2},q_{n_2}}p_{n_2}^{n_2-k_2+1}}{[n_2+1]_{p_{n_2},q_{n_2}}}$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c}n_1\\k_1\end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c}n_2\\k_2\end{array}\right]_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}$$

$$=\tfrac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\tfrac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}\textstyle\sum_{k_1=1}^{n_1}p_{n_1}^{n_1-k_1+1+\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c}n_1-1\\k_1-1\end{array}\right]_{p_{n_1},q_{n_1}}(q_{n_1}x)^{k_1}$$

$$=\tfrac{x}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\tfrac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}\textstyle\sum_{k_1=0}^{n_1}p_{n_1-1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c}n_1-1\\k_1\end{array}\right]_{p_{n_1},q_{n_1}}(q_{n_1}x)^{k_1}$$

$= \frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}\frac{x}{1+x}.$

(iii) It can be proved in a similar way.

(iv) $L_{p,q}^{n}\left(e_{20};x,y\right) = \frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_1=1}^{n_1}\sum_{k_2=1}^{n_2}\frac{[k_1]^2_{p_{n_1},q_{n_1}}p_{n_1}^{2(n_1-k_1+1)}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}$
$\times p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c} n_1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c} n_2 \\ k_2 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}y^{k_2}.$

Using (1.6), we get

$$L_{p,q}^{n}\left(\frac{u^2}{(1+u)^2};x\right)$$
$$= \frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\sum_{k_1=2}^{n_1}\frac{q_{n_1}[k_1]_{p_{n_1},q_{n_1}}[k_1-1]_{p_{n_1},q_{n_1}}p_{n_1}^{2n_1-2k_1+2}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}$$
$$+\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\sum_{k_1=1}^{n_1}p_{n_1}^{k_1-1}\frac{[k_1]_{p_{n_1},q_{n_1}}p_{n_1}^{2n_1-2k_1+2}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}$$

$$= \frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{q_{n_1}[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\sum_{k_1=2}^{n_1}p_{n_1}^{\left((2n_1-2k_1+2)+\frac{(n_1-k_1)(n_1-k_1-1)}{2}\right)}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1-2 \\ k_1-2 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}$$

$$+\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\sum_{k_1=1}^{n_1}p_{n_1}^{\left((k_1-1)+(2n_1-2k_1+2)+\frac{(n_1-k_1)(n_1-k_1-1)}{2}\right)}q_{n_{n_1}}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1-1 \\ k_1-1 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}$$

$$= x^2\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{q_{n_1}[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\sum_{k_1=0}^{n_1-2}p_{n_1}^{\left((2n_1-2k_1-2)+\frac{(n_1-k_1-2)(n_1-k_1-3)}{2}\right)}q_{n_1}^{\frac{(k_1+1)(k_1+2)}{2}}\left[\begin{array}{c} n_1-2 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}$$

$$+x\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\sum_{k_1=0}^{n_1-1}p_{n_1}^{\left(k_1+(2n_1-2k_1)+\frac{(n_1-k_1-1)(n_1-k_1-2)}{2}\right)}q_{n_1}^{\frac{k_1(k_1+1)}{2}}\left[\begin{array}{c} n_1-1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}x^{k_1}$$

$$= x^2\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{p_{n_1}q_{n_1}^2[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\sum_{k_1=0}^{n_1-2}p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1-2 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}(q_{n_1}^2x)^{k_1}$$

$$+x\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{p_{n_1}^{n_1+1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\sum_{k_1=0}^{n_1-1}p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1-1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}(q_{n_1}x)^{k_1}$$

$$= \frac{p_{n_1}q_{n_1}^2[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\frac{x^2}{(1+x)(p_{n_1}+q_{n_1}x)}+\frac{p_{n_1}^{n_1+1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}\left(\frac{x}{1+x}\right).$$

(v) It can be proved in a similar way.

□

Lemma 2.2. *The operators defined in (2.1) satisfy the following conditions.*

(i) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y) = A_{n_1}^{(p_{n_1},q_{n_1}),x}\left(B_{n_2}^{(p_{n_2},q_{n_2}),y}(f;x,y)\right)$

(ii) $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y) = B_{n_2}^{(p_{n_2},q_{n_2}),y}\left(A_{n_1}^{(p_{n_1},q_{n_1}),x}(f;x,y)\right)$

Proof. (i) $A_{n_1}^{(p_{n_1},q_{n_1}),x}\left(B_{n_2}^{(p_{n_2},q_{n_2}),y}(f;x,y)\right)$

$$
\begin{aligned}
&= A_{n_1}^{(p_{n_1},q_{n_1}),x}\left(\frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_2=0}^{n_2} f\left(x,\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right)p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c} n_2 \\ k_2 \end{array}\right]_{p_{n_2},q_{n_2}} y^{k_2}\right)\\
&= \frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_2=0}^{n_2} A_{n_1}^{(p_{n_1},q_{n_1}),x}\left(f\left(x,\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right)\right)p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c} n_2 \\ k_2 \end{array}\right]_{p_{n_2},q_{n_2}} y^{k_2}\\
&= \frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_2=0}^{n_2} p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c} n_2 \\ k_2 \end{array}\right]_{p_{n_2},q_{n_2}} y^{k_2}\\
&\times \sum_{k_1=0}^{n_1}\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)} f\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{[n_1-k_1+1]_{p_{n_1},q_{n_1}}q_{n_1}^{k_1}},\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right)\\
&\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}\left[\begin{array}{c} n_1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}} x^{k_1}\\
&= \frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2} f\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{[n_1-k_1+1]_{p_{n_1},q_{n_1}}q_{n_1}^{k_1}},\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right)\\
&\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c} n_1 \\ k_1 \end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c} n_2 \\ k_2 \end{array}\right]_{p_{n_2},q_{n_2}} x^{k_1}y^{k_2}\\
&= L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y).
\end{aligned}
$$

(ii) It can be proved in a similar way.

□

3. APPROXIMATION PROPERTIES OF BIVARIATE OPERATORS

In this section, we obtain the Korovkin's type approximation theorem for our operators defined in (2.1).

Let $C_{\mathcal{B}}(\mathbb{R}_+^2)$ be the set of all bounded and continuous functions on $\mathbb{R}_+^2$ which is linear normed space with

$$\| f \|_{C_B(\mathbb{R}_+^2)} = \sup_{x,y\geq 0} \mid f(x,y) \mid .$$

If

$$\lim_{n,m\to\infty} \| f_{n,m} - f \|_{C_{\mathcal{B}}(\mathbb{R}_+^2)} = 0$$

holds, then we say that the sequence $\{f_{n,m}\}$ converges uniformly to f and it is written as $f_{n,m} \rightrightarrows f$.

Now, let us introduce modulus of continuity $\omega(\delta)$ which satisfy the following conditions:

(i) $\omega(\delta)$ is a non-negative increasing function for δ on $\mathbb{R}_+$
(ii) $\omega(\delta_1+\delta_2)\leq \omega(\delta_1)+\omega(\delta_2)$
(iii) $\lim_{\delta\to 0}\omega(\delta)=0$.

Let H_ω be the space of all real-valued functions f satisfying the condition

$$\mid f(x) - f(y) \mid \leq \omega\left(\left|\frac{x}{1+x}-\frac{y}{1+y}\right|\right),$$

for any $x,y\in\mathbb{R}_+$.
Note that $H_\omega \subset C_{\mathcal{B}}(\mathbb{R}_+)$ for the bounded and continuous function f on $\mathbb{R}_+$.

Theorem 3.1 ([11]). *Let $\{A_n\}$ be the sequence of positive linear operators from $H_\omega(\mathbb{R}_+)$ into $C_B(\mathbb{R}_+)$, satisfying the conditions*

$$\lim_{n\to\infty} \| A_n\left(\left(\frac{t}{1+t}\right)^\nu ; x\right) - \left(\frac{x}{1+x}\right)^\nu \|_{C_B},$$

for $\nu = 0, 1, 2$. Then for any function $f \in H_\omega$

$$\lim_{n\to\infty} \| A_n(f) - f \|_{C_B} = 0.$$

In order to get the convergence results for the operators $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$, we take $p = p_{n_1},\ p = p_{n_2}$ and $q = q_{n_1},\ q = q_{n_2}$ where $q_{n_i} \in (0,1)$ and $p_{n_i} \in (q_{n_i}, 1]$ for $i = 1, 2$ satisfying,

$$\lim_{n_1,n_2\to\infty} p_{n_1},\ p_{n_2} \to 1, \text{ and } \lim_{n_1,n_2\to\infty} q_{n_1},\ q_{n_2} \to 1 \tag{3.1}$$

Now we prove the bivariate case of Theorem 3.1.

Theorem 3.2. *Let $p = p_{n_1},\ p = p_{n_2}$ and $q = q_{n_1},\ q = q_{n_2}$ satisfying (3.1), for $0 < q_{n_1} < p_{n_1} \leq 1,\ 0 < q_{n_2} < p_{n_2} \leq 1$ and if $A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$ be the sequence of a linear positive operator, defined as $A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})} : H_\omega(\mathbb{R}_+^2) \to C_B(\mathbb{R}_+^2)$, satisfying the following conditions:*

(i) $\lim_{n_1,n_2\to\infty} \| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00}; x, y) - e_{00} \|_{C_{\mathbb{R}_+^2}} = 0,$

(ii) $\lim_{n_1,n_2\to\infty} \| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{10}; x, y) - e_{10} \|_{C_{\mathbb{R}_+^2}} = 0,$

(iii) $\lim_{n_1,n_2\to\infty} \| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{01}; x, y) - e_{01} \|_{C_{\mathbb{R}_+^2}} = 0,$

(iv) $\lim_{n_1,n_2\to\infty} \| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{20}; x, y) - e_{20} \|_{C_{\mathbb{R}_+^2}} = 0,$

(v) $\lim_{n_1,n_2\to\infty} \| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{02}; x, y) - e_{02} \|_{C_{\mathbb{R}_+^2}} = 0.$

Then for any function $f \in H_\omega(\mathbb{R}_+^2)$

$$\lim_{n_1,n_2\to\infty} \| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f; x, y) - f(x, y) \|_{C_{\mathbb{R}_+^2}} = 0 \tag{3.2}$$

holds. Here $e_{ij} : \mathbb{R}_+^2 \to [0, 1)$ defined in (1.7) are two dimensional test functions.

Proof. For the bivariate case on $H_\omega(\mathbb{R}_+^2)$, we have

$$| f(u, v) - f(x, y) | \leq \omega \left(\left| \left(\frac{u}{1+u}, \frac{v}{1+v} \right) - \left(\frac{x}{1+x}, \frac{y}{1+y} \right) \right| \right).$$

If $f \in H_\omega(\mathbb{R}_+^2)$, then for any $\epsilon > 0$, there exits $\delta > 0$ such that

$$| f(u, v) - f(x, y) | < \epsilon.$$

Also if

$$\sqrt{\left(\frac{u}{1+u} - \frac{x}{1+x} \right)^2 + \left(\frac{v}{1+v} - \frac{y}{1+y} \right)^2} < \delta$$

boundedness of f implies that there exits a positive constant M such that

$$| f(u, v) - f(x, y) | \leq \frac{2M}{\delta^2} \left[\left(\frac{u}{1+u} - \frac{x}{1+x} \right)^2 + \left(\frac{v}{1+v} - \frac{y}{1+y} \right)^2 \right],$$

if $\sqrt{\left(\frac{u}{1+u} - \frac{x}{1+x} \right)^2 + \left(\frac{v}{1+v} - \frac{y}{1+y} \right)^2} \geq \delta.$

Therefore, for all $(u, v),\ (x, y) \in \mathbb{R}_+^2$, we have

$$| f(u, v) - f(x, y) | \leq \epsilon + \frac{2M}{\delta^2} \left[\left(\frac{u}{1+u} - \frac{x}{1+x} \right)^2 + \left(\frac{v}{1+v} - \frac{y}{1+y} \right)^2 \right].$$

Now applying the operator $A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$ to the above inequality, we get

$| A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f; x, y) - f |_{C_{\mathbb{R}_+^2}}$

$\leq (\epsilon + M) \mid A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y) - e_{00} \mid + \epsilon$
$+\frac{2M}{\delta^2} \mid A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{20}+e_{02};x,y) - (e_{20}+e_{02}) \mid$
$+\frac{4M}{\delta^2} \mid A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{10};x,y) - e_{10} \mid + \frac{4M}{\delta^2} \mid A_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{01};x,y) - e_{01} \mid .$
By using the conditions (i)-(v) we get the result. □

Theorem 3.3. *Let* $p = p_{n_1},\ p = p_{n_2}$ *and* $q = q_{n_1},\ q = q_{n_2}$ *satisfying (3.1), for* $0 < q_{n_1} < p_{n_1} \leq 1,\ 0 < q_{n_2} < p_{n_2} \leq 1$ *and if* $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$ *be the sequence of a linear positive operator defined in (2.1) such that* $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})} : H_\omega(\mathbb{R}_+^2) \to C_B(\mathbb{R}_+^2)$ *, satisfying the conditions of Theorem 3.2. Then for any* $f \in H_\omega(\mathbb{R}_+^2)$*,* $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$ *converges uniformly to* f*. That is* $\forall f \in H_\omega(\mathbb{R}_+^2)$*, we have*

$$\lim_{n_1,n_2\to\infty} \parallel L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y) - f(x,y) \parallel_{C_{\mathbb{R}_+^2}} = 0.$$

Proof. Using Theorem 3.1 we see that it is sufficient to verify the following conditions:

$$\lim_{n_1,n_2\to\infty} \parallel L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{ij};x,y) - e_{ij} \parallel_{C_{\mathbb{R}_+^2}} = 0,\ i,j = 0,1,2, \tag{3.3}$$

here e_{ij} are the test functions defined in (1.7). From Lemma 2.1, the first condition of (3.3) is fulfilled for $i,j = 0$. Now it is easy to see that from (ii) of Lemma 2.1

$$\begin{aligned}\lim_{n_1,n_2\to\infty} \parallel L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{10};x,y) - e_{10} \parallel_{C_{\mathbb{R}_+^2}} &= \sup_{x,y\geq 0} \left| \frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}} \frac{x}{1+x} - \frac{x}{1+x} \right| \\ &\leq \left| \frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}} - 1 \right| \\ &\leq \left| \left(\frac{p_{n_1}}{q_{n_1}}\right) \left(1 - p_{n_1}^{n_1} \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}}\right) - 1 \right|\end{aligned}$$

since $q_{n_1}[n_1]_{p_{n_1},q_{n_1}} = [n_1+1]_{p_{n_1},q_{n_1}} - p_{n_1}^{n_1},\ [n_1+1]_{p_{n_1},q_{n_1}} \to \infty$ as $n_1 \to \infty$, the condition (3.3) holds for $i = 1,\ j = 0$.
Similarly for $i = 0,\ j = 1$ it can be shown easily.

To verify this condition for $i = 2,\ j = 0$, consider (iv) of Lemma 2.1. Then we see that
$\lim_{n_1\to\infty} \parallel L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{20};x,y) - e_{20} \parallel_{C_{\mathbb{R}_+^2}}$
$= \sup_{x\geq 0} \left\{ \frac{x^2}{(1+x)^2} \left(\frac{p_{n_1} q_{n_1}^2 [n_1]_{p_{n_1},q_{n_1}} [n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2} . \frac{1+x}{p_{n_1}+q_{n_1}x} - 1 \right) + \frac{p_{n_1}^{n_1+1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2} . \frac{x}{1+x} \right\}.$
A small calculation leads to

$$\frac{[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2} = \frac{1}{q_{n_1}^3} \left\{ 1 - p_{n_1}^{n_1} \left(2 + \frac{q_{n_1}}{p_{n_1}}\right) \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}} + (p_{n_1}^{n_1})^2 \left(1 + \frac{q_{n_1}}{p_{n_1}}\right) \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}^2} \right\},$$

and

$$\frac{[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2} = \frac{1}{q_{n_1}} \left(\frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}} - p_{n_1}^{n_1} \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}^2} \right).$$

Thus we have

$\lim_{n_1\to\infty} \parallel L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{20};x,y) - e_{20} \parallel_{C_{\mathbb{R}_+^2}}$

$\leq \frac{p_{n_1}}{q_{n_1}} \left\{ 1 - p_{n_1}^{n_1} \left(2 + \frac{q_{n_1}}{p_{n_1}}\right) \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}} + (p_{n_1}^{n_1})^2 \left(1 + \frac{q_{n_1}}{p_{n_1}}\right) \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}^2} - 1 \right\}$
$+ p_{n_1}^{n_1} . \frac{p_{n_1}}{q_{n_1}} \left(\frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}} - p_{n_1}^{n_1} \frac{1}{[n_1+1]_{p_{n_1},q_{n_1}}^2} \right).$

This implies that the condition (3.3) holds for also $i = 2,\ j = 0$ and also in similar way for $i = 0,\ j = 2$ the proof is true. This completes the proof. $\square$

4. RATE OF CONVERGENCE

In this section, rate of convergence of operators (2.1) by means of modulus of continuity of some bivariate modulus of smoothness functions are introduced.
The modulus of continuity for bivariate case is defined as follows. For $f \in H_\omega(\mathbb{R}_+^2)$ is defined by

$$\widetilde{\omega}(f;\delta_1,\delta_2) = \sup\nolimits_{u,x\geq 0}\left\{\Big|f(u,v)-f(x,y)\Big|;\ \left|\tfrac{u}{1+u}-\tfrac{x}{1+x}\right| \leq \delta_1, \left|\tfrac{v}{1+v}-\tfrac{y}{1+y}\right| \leq \delta_2,\ (u,v)\in\mathbb{R}_+^2,\ (x,y)\in\mathbb{R}_+^2\right\}.$$

where for all $f \in H_\omega(\mathbb{R}_+)$ $\widetilde{\omega}(f;\delta_1,\delta_2)$ satisfies the following conditions:

(i) $\lim_{\delta_1,\delta_2\to 0}\widetilde{\omega}(f;\delta_1,\delta_2) \to 0$

(ii) $\mid f(u,v)-f(x,y)\mid \leq \widetilde{\omega}(f;\delta_1,\delta_2)\left(\frac{\left|\frac{u}{1+u}-\frac{x}{1+x}\right|}{\delta_1}+1\right)\left(\frac{\left|\frac{v}{1+v}-\frac{y}{1+y}\right|}{\delta_2}+1\right)$.

Theorem 4.1. *Let $p = p_{n_1},\ p = p_{n_2}$ and $q = q_{n_1},\ q = q_{n_2}$ satisfying (3.1), for $0 < q_{n_1} < p_{n_1} \leq 1,\ 0 < q_{n_2} < p_{n_2} \leq 1$ and if $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$ be the sequence of a linear positive operator defined in (2.1). Then for all $f \in H_\omega(\mathbb{R}_+^2)$ and $x, y \geq 0$, we have*

$$\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y)-f(x,y)\mid \leq 4\omega(f;\delta_{n_1}(x),\delta_{n_2}(y)) \tag{4.1}$$

where

$$\delta_{n_1}(x) = \frac{x^2}{(1+x)^2}\left(\frac{p_{n_1}q_{n_1}^2[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2}\frac{1+x}{p_{n_1}+q_{n_1}x}-2\frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}+1\right)+\frac{p_{n_1}^{n_1+1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2}\frac{x}{1+x}$$

$$\delta_{n_2}(y) = \frac{y^2}{(1+y)^2}\left(\frac{p_{n_2}q_{n_2}^2[n_2]_{p_{n_2},q_{n_2}}[n_2-1]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}^2}\frac{1+y}{p_{n_2}+q_{n_2}y}-2\frac{p_{n_2}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}+1\right)+\frac{p_{n_2}^{n_2+1}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}^2}\frac{y}{1+y}.$$

Proof.

$$\begin{aligned}
\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y)-f(x,y)\mid &\leq L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(\mid f(u,v)-f(x,y)\mid;x,y)\\
&\leq \widetilde{\omega}(f;\delta_{n_1},\delta_{n_2})\left\{1+\frac{1}{\delta_{n_1}}L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\Big|\frac{u}{1+u}-\frac{x}{1+x}\Big|;x,y\right)\right\}\\
&\times \left\{1+\frac{1}{\delta_{n_2}}L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\Big|\frac{v}{1+v}-\frac{y}{1+y}\Big|;x,y\right)\right\}.
\end{aligned}$$

Now by using the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
&\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y)-f(x,y)\mid\\
&\leq \widetilde{\omega}(f;\delta_{n_1},\delta_{n_2})\left\{1+\frac{1}{\delta_{n_1}}\left[\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\frac{u}{1+u}-\frac{x}{1+x}\right)^2;x,y\right)\right]^{\frac{1}{2}}\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)\right)^{\frac{1}{2}}\right\}\\
&\times \widetilde{\omega}(f;\delta_{n_1},\delta_{n_2})\left\{1+\frac{1}{\delta_{n_2}}\left[\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\frac{v}{1+v}-\frac{y}{1+y}\right)^2;x,y\right)\right]^{\frac{1}{2}}\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)\right)^{\frac{1}{2}}\right\}
\end{aligned}$$

$$\begin{aligned}
&\leq \widetilde{\omega}(f;\delta_{n_1},\delta_{n_2})\left\{1+\tfrac{1}{\delta_{n_1}}\left[\tfrac{x^2}{(1+x)^2}\left(\tfrac{p_{n_1}q_{n_1}^2[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2}\tfrac{1+x}{p_{n_1}+q_{n_1}x}-2\tfrac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}+1\right)+\tfrac{p_{n_1}^{n_1+1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}^2}\tfrac{x}{1+x}\right]^{\frac{1}{2}}\right.\\
&\times\left\{1+\tfrac{1}{\delta_{n_2}}\left[\tfrac{y^2}{(1+y)^2}\left(\tfrac{p_{n_2}q_{n_2}^2[n_2]_{p_{n_2},q_{n_2}}[n_2-1]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}^2}\tfrac{1+y}{p_{n_2}+q_{n_2}y}-2\tfrac{p_{n_2}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}+1\right)+\tfrac{p_{n_2}^{n_2+1}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}^2}\tfrac{y}{1+y}\right]^{\frac{1}{2}}\right\}.
\end{aligned}$$

This completes the proof. $\square$

Now we will give an estimate concerning the rate of convergence by means of Lipschtz type maximal functions. In [8, 9], the Lipschtz type maximal function space on $E \times E \subset \mathbb{R}_+ \times \mathbb{R}_+$ is defined as follows:

$$\widetilde{W}_{\alpha_1,\alpha_2 E^2} = \{f : \sup(1+u)^{\alpha_1}(1+v)^{\alpha_2}\left(f_{\alpha_1,\alpha_2}(u,v) - f_{\alpha_1,\alpha_2}(x,y)\right)$$

$$\leq M \frac{1}{(1+x)^{\alpha_1}} \frac{1}{(1+y)^{\alpha_2}} : x, y \leq 0, \text{ and } (u,v) \in E^2\}$$

where f is bounded and continuous function on $\mathbb{R}_+$, M is a positive constant. For $0 \leq \alpha_1 \leq 1,\ 0 \leq \alpha_2 \leq 1$, we define the function f_{α_1,α_2} as follows:

$$f_{\alpha_1,\alpha_2}(u,v) - f_{\alpha_1,\alpha_2}(x,y) = \frac{\mid f(u,v) - f(x,y) \mid}{\mid u - x \mid^{\alpha_1} \mid v - y \mid^{\alpha_2}}$$

Also, let $d(x,E)$ be the distance between x and E, that is

$$d(x,E) = \inf\{\mid x - y \mid; y \in E\}.$$

Theorem 4.2. *Let $0 < \alpha_1 \leq 1,\ 0 < \alpha_2 \leq 1$, then for all $f \in \widetilde{W}_{\alpha_1,\alpha_2,E^2}$ we have*

$$\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y) - f(x,y) \mid$$

$$\leq M\left(\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y) + \delta_{n_1}^{\frac{\alpha_1}{2}}(x)\left(d(x,E)\right)^{\alpha_1} + \delta_{n_2}^{\frac{\alpha_2}{2}}(y)\left(d(y,E)\right)^{\alpha_2} + 2\left(d(x,E)\right)^{\alpha_1}\left(d(y,E)\right)^{\alpha_2}\right)$$

where $\delta_{n_1}(x)$ and $\delta_{n_2}(y)$ are defined in Theorem 4.1.

Proof. Let $x, y \in \mathbb{R}_+$, then there exits a $(x_0, y_0) \in E \times E$ such that $\mid x - x_0 \mid = d(x,E)$, and $\mid y - y_0 \mid = d(y,E)$. Thus we can write

$$\mid f(u,v) - f(x,y) \mid \leq \mid f(u,v) - f(x_0,y_0) \mid + \mid f(x_0,y_0) - f(x,y) \mid.$$

Since $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$ is a positive linear operator, $f \in \widetilde{W}_{\alpha_1,\alpha_2,E^2}$ and by using the previous inequality we have

$$\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y) - f(x,y) \mid$$
$$\leq \mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(\mid f(u,v) - f(x_0,y_0) \mid; x,y) + \mid f(x_0,y_0) - f(x,y) \mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)$$

$$\leq M L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\tfrac{u}{1+u} - \tfrac{x_0}{1+x_0}\right|^{\alpha_1}\left|\tfrac{v}{1+v} - \tfrac{y_0}{1+y_0}\right|^{\alpha_2}; x, y\right)$$
$$+ M\left|\tfrac{x}{1+x} - \tfrac{x_0}{1+x_0}\right|^{\alpha_1}\left|\tfrac{y}{1+y} - \tfrac{y_0}{1+y_0}\right|^{\alpha_2} L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)$$

Since $(a+b)^\alpha \leq a^\alpha + b^\alpha$, for $a, b \geq 0$ and $0 \leq \alpha \leq 1$, which consequently imply

$$\left|\frac{u}{1+u} - \frac{x_0}{1+x_0}\right|^{\alpha_1} \leq \left|\frac{u}{1+u} - \frac{x}{1+x}\right|^{\alpha_1} + \left|\frac{x}{1+x} - \frac{x_0}{1+x_0}\right|^{\alpha_1}$$

$$\left|\frac{v}{1+v} - \frac{y_0}{1+y_0}\right|^{\alpha_2} \leq \left|\frac{v}{1+v} - \frac{y}{1+y}\right|^{\alpha_2} + \left|\frac{y}{1+y} - \frac{y_0}{1+y_0}\right|^{\alpha_2}$$

$$
\begin{aligned}
\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y)-f(x,y)\mid &\leq L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\frac{u}{1+u}-\frac{x}{1+x}\right|^{\alpha_1}\left|\frac{v}{1+v}-\frac{y}{1+y}\right|^{\alpha_2};x,y\right)\\
&+\left|\frac{y}{1+y}-\frac{y_0}{1+y_0}\right|^{\alpha_2}L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\frac{u}{1+u}-\frac{x}{1+x}\right|^{\alpha_1};x,y\right)\\
&+\left|\frac{x}{1+x}-\frac{x_0}{1+x_0}\right|^{\alpha_1}L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\frac{v}{1+v}-\frac{y}{1+y}\right|^{\alpha_2};x,y\right)\\
&+\left|\frac{x}{1+x}-\frac{x_0}{1+x_0}\right|^{\alpha_1}\left|\frac{y}{1+y}-\frac{y_0}{1+y_0}\right|^{\alpha_2}L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y).
\end{aligned}
$$

By applying the Hölder inequality with $p_1=\frac{2}{\alpha_1},\ p_2=\frac{2}{\alpha_2}$ and $q_1=\frac{2}{2-\alpha_1},\ q_2=\frac{2}{2-\alpha_2}$, then we have

$$
\begin{aligned}
&L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\tfrac{u}{1+u}-\tfrac{x}{1+x}\right|^{\alpha_1}\left|\tfrac{v}{1+v}-\tfrac{y}{1+y}\right|^{\alpha_2};x,y\right)\\
&=L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\tfrac{u}{1+u}-\tfrac{x}{1+x}\right|^{\alpha_1};x,y\right)L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\tfrac{v}{1+v}-\tfrac{y}{1+y}\right|^{\alpha_2};x,y\right)\\
&\leq\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\tfrac{u}{1+u}-\tfrac{x}{1+x}\right|^{2};x,y\right)\right)^{\frac{\alpha_1}{2}}\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)\right)^{\frac{2-\alpha_1}{2}}\\
&\times\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}\left(\left|\tfrac{v}{1+v}-\tfrac{y}{1+y}\right|^{2};x,y\right)\right)^{\frac{\alpha_2}{2}}\left(L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(e_{00};x,y)\right)^{\frac{2-\alpha_2}{2}}.
\end{aligned}
$$

This completes the proof. □

Corollary 4.3. *If it is taken $E=\mathbb{R}_+$ as a special case of Theorem 4.2, then for all $f\in\widetilde{W}_{\alpha_1,\alpha_2,\mathbb{R}_+^2}$, we have*

$$
\mid L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}(f;x,y)-f(x,y)\mid\leq\mathcal{M}\left(\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y)\right),
$$

where $\delta_{n_1}(x)$ and $\delta_{n_2}(y)$ are defined in Theorem 4.1.

5. SOME GENERALIZATION OF $L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}$

In this section, similar as in [4, 7], we define some generalization of the operators (2.1) for (p,q)-integers as follows:

$$
\begin{aligned}
&L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}(f;x,y)\\
&=\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}f\left(\frac{p_{n_1}^{n_1+1-k_1}[k_1]_{p_{n_1},q_{n_1}}+\gamma_1}{b_{n_1,k_1}},\frac{p_{n_2}^{n_2+1-k_2}[k_2]_{p_{n_2},q_{n_2}}+\gamma_2}{b_{n_2,k_2}}\right)\\
&\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c}n_1\\k_1\end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c}n_2\\k_2\end{array}\right]_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2},\ (\gamma_1,\gamma_2\in\mathbb{R})
\end{aligned}
\tag{5.1}
$$

where b_{n_i,k_i} for $i=1,2$ satisfies the following conditions:

$$
p_{n_i}^{n_i-k_i+1}[k_i]_{p_{n_i},q_{n_i}}+b_{n_i,k_i}=c_{n_i}\text{ and }\frac{[n_i]_{p_{n_i},q_{n_i}}}{c_{n_i}}\to 1,\ \text{for } n_i\to\infty.
$$

It is easy to check that if $b_{n_i,k_i}=q_{n_i}^{k_i}[n_i-k_i+1]_{p_{n_i},q_{n_i}}+\beta_i$ for any n_i,k_i and $0<q_{n_i}<p_{n_i}\leq 1$, then $c_{n_i}=[n_i+1]_{p_{n_i},q_{n_i}}+\beta_i$. If we choose $p_{n_i}=1,\ \gamma_i=0$, for $i=1,2$ then operators reduces to q-bivariate-BBH opeartors defined in [8, 9].

Theorem 5.1. *Let* $p = p_{n_1},\ p = p_{n_2}$ *and* $q = q_{n_1},\ q = q_{n_2}$ *satisfying (3.1), for* $0 < q_{n_1} < p_{n_1} \le 1,\ 0 < q_{n_2} < p_{n_2} \le 1$ *and if* $L^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}_{n_1,n_2}$ *is defined by (5.1), then for any function* $f \in \widetilde{W}_{\alpha_1,\alpha_2,E^2}$*, we have*

$$\lim_{n_1,n_2} \parallel L^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}_{n_1,n_2}(f;x,y) - f(x,y) \parallel_{C_B(\mathbb{R}^2_+)}$$

$$\le 3M \max\left(\frac{[n_1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right)^{\alpha_1}\left(\frac{\gamma_1}{[n_1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left(\frac{[n_2]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right)^{\alpha_2}\left(\frac{\gamma_2}{[n_2]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2},$$

$$\left|1-\frac{[n_1+1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right|^{\alpha_1}\left(\frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left|1-\frac{[n_2+1]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right|^{\alpha_2}\left(\frac{p_{n_2}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2},$$

$$\left(1-2\frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}+\frac{[n_1]_{p_{n_1},q_{n_1}}[n_1-1]_{p_{n_1},q_{n_1}}}{[n_1+1]^2_{p_{n_1},q_{n_1}}}q_{n_1}\right)\left(1-2\frac{p_{n_2}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}+\frac{[n_2]_{p_{n_2},q_{n_2}}[n_2-1]_{p_{n_2},q_{n_2}}}{[n_2+1]^2_{p_{n_2},q_{n_2}}}q_{n_2}\right).$$

Proof. Using (2.1) and (5.1) we have

$$\left|L^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}_{n_1,n_2}(f;x,y) - f(x,y)\right| \le \frac{1}{\ell^{(p_{n_1},q_{n_1})}_{n_1}(x)}\frac{1}{\ell^{(p_{n_2},q_{n_2})}_{n_2}(y)}\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}$$

$$\times\left|f\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}+\gamma_1}{b_{n_1,k_1}},\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}+\gamma_2}{b_{n_2,k_2}}\right)-f\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{\gamma_1+b_{n_1,k_1}},\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{\gamma_2+b_{n_2,k_2}}\right)\right|$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\begin{bmatrix} n_1 \\ k_1 \end{bmatrix}_{p_{n_1},q_{n_1}}\begin{bmatrix} n_2 \\ k_2 \end{bmatrix}_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}$$

$$+\frac{1}{\ell^{(p_{n_1},q_{n_1})}_{n_1}(x)}\frac{1}{\ell^{(p_{n_2},q_{n_2})}_{n_2}(y)}\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}$$

$$\times\left|f\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{\gamma_1+b_{n_1,k_1}},\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{\gamma_2+b_{n_2,k_2}}\right)-f\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{[n_1-k_1+1]_{p_{n_1},q_{n_1}}q_{n_1}^{k_1}},\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right)\right|$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\begin{bmatrix} n_1 \\ k_1 \end{bmatrix}_{p_{n_1},q_{n_1}}\begin{bmatrix} n_2 \\ k_2 \end{bmatrix}_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}$$

$$+\mid L^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2})}_{n_1,n_2}(f;x,y) - f(x,y)\mid.$$

Since $f \in \widetilde{W}_{\alpha_1,\alpha_2,E^2}$ and by using the Corollary 4.3 we can write

$$\left|L^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}_{n_1,n_2}(f;x,y) - f(x,y)\right|$$

$$\le M\frac{1}{\ell^{(p_{n_1},q_{n_1})}_{n_1}(x)}\frac{1}{\ell^{(p_{n_2},q_{n_2})}_{n_2}(y)}\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\left|\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}+\gamma_1}{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}+\gamma_1+b_{n_1,k_1}}-\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{\gamma_1+p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}+b_{n_1,k_1}}\right|^{\alpha_1}$$

$$\times\left|\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}+\gamma_2}{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}+\gamma_2+b_{n_2,k_2}}-\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{\gamma_2+p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}+b_{n_2,k_2}}\right|^{\alpha_2}$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\begin{bmatrix} n_1 \\ k_1 \end{bmatrix}_{p_{n_1},q_{n_1}}\begin{bmatrix} n_2 \\ k_2 \end{bmatrix}_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}$$

$$+M\frac{1}{\ell^{(p_{n_1},q_{n_1})}_{n_1}(x)}\frac{1}{\ell^{(p_{n_2},q_{n_2})}_{n_2}(y)}\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\left|\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}+\gamma_1+b_{n_1,k_1}}-\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}+[n_1-k_1+1]_{p_{n_1},q_{n_1}}q_{n_1}^{k_1}}\right|$$

$$\times\left|\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}+\gamma_2+b_{n_2,k_2}}-\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}+[n_2-k_2+1]_{p_{n_2},q_{n_2}}q_{n_2}^{k_2}}\right|$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\begin{bmatrix} n_1 \\ k_1 \end{bmatrix}_{p_{n_1},q_{n_1}}\begin{bmatrix} n_2 \\ k_2 \end{bmatrix}_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}$$

$$+M\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y).$$

This implies that

$$\left|L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}(f;x,y)-f(x,y)\right|$$

$$\leq M\left(\frac{[n_1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right)^{\alpha_1}\left(\frac{\gamma_1}{[n_1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left(\frac{[n_2]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right)^{\alpha_2}\left(\frac{\gamma_2}{[n_2]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2}$$

$$+M\frac{1}{\ell_{n_1}^{(p_{n_1},q_{n_1})}(x)}\frac{1}{\ell_{n_2}^{(p_{n_2},q_{n_2})}(y)}\left|1-\frac{[n_1+1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right|^{\alpha_1}\left|1-\frac{[n_2+1]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right|^{\alpha_2}$$

$$\times\sum_{k_1=0}^{n_1}\sum_{k_2=0}^{n_2}\left(\frac{p_{n_1}^{n_1-k_1+1}[k_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left(\frac{p_{n_2}^{n_2-k_2+1}[k_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2}$$

$$\times p_{n_1}^{\frac{(n_1-k_1)(n_1-k_1-1)}{2}}q_{n_1}^{\frac{k_1(k_1-1)}{2}}p_{n_2}^{\frac{(n_2-k_2)(n_2-k_2-1)}{2}}q_{n_2}^{\frac{k_2(k_2-1)}{2}}\left[\begin{array}{c}n_1\\k_1\end{array}\right]_{p_{n_1},q_{n_1}}\left[\begin{array}{c}n_2\\k_2\end{array}\right]_{p_{n_2},q_{n_2}}x^{k_1}y^{k_2}+M\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y)$$

$$=M\left(\frac{[n_1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right)^{\alpha_1}\left(\frac{\gamma_1}{[n_1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left(\frac{[n_2]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right)^{\alpha_2}\left(\frac{\gamma_2}{[n_2]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2}$$

$$+M\left|1-\frac{[n_1+1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right|^{\alpha_1}\left|1-\frac{[n_2+1]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right|^{\alpha_2}L_{p_{n_1},q_{n_1},p_{n_2},q_{n_2}}^{n_1,n_2}\left(\left(\frac{u}{1+u}\right)^{\alpha_1};x,y\right)L_{p_{n_1},q_{n_1},p_{n_2},q_{n_2}}^{n_1,n_2}\left(\left(\frac{v}{1+v}\right)^{\alpha_2};x,y\right)$$

$$+M\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y).$$

Using the Hölder inequality for $p_1=\frac{1}{\alpha_1},\ q_1=\frac{1}{1-\alpha_1}$, and $p_2=\frac{1}{\alpha_2},\ q_2=\frac{1}{1-\alpha_2}$ we get

$$\left|L_{n_1,n_2}^{(p_{n_1},p_{n_2}),(q_{n_1},q_{n_2}),(\gamma_1,\gamma_2)}(f;x,y)-f(x,y)\right|$$

$$\leq M\left(\frac{[n_1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right)^{\alpha_1}\left(\frac{\gamma_1}{[n_1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left(\frac{[n_2]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right)^{\alpha_2}\left(\frac{\gamma_2}{[n_2]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2}$$

$$+M\left|1-\frac{[n_1+1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right|^{\alpha_1}\left|1-\frac{[n_2+1]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right|^{\alpha_2}L_{p_{n_1},q_{n_1},p_{n_2},q_{n_2}}^{n_1,n_2}\left(\frac{u}{1+u};x,y\right)^{\alpha_1}\left(L_{p_{n_1},q_{n_1},p_{n_2},q_{n_2}}^{n_1,n_2}(e_{00};x,y)\right)^{1-\alpha_1}$$

$$\times L_{p_{n_1},q_{n_1},p_{n_2},q_{n_2}}^{n_1,n_2}\left(\frac{v}{1+v};x,y\right)^{\alpha_2}\left(L_{p_{n_1},q_{n_1},p_{n_2},q_{n_2}}^{n_1,n_2}(e_{00};x,y)\right)^{1-\alpha_2}+M\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y)$$

$$\leq M\left(\frac{[n_1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right)^{\alpha_1}\left(\frac{\gamma_1}{[n_1]_{p_{n_1},q_{n_1}}}\right)^{\alpha_1}\left(\frac{[n_2]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right)^{\alpha_2}\left(\frac{\gamma_2}{[n_2]_{p_{n_2},q_{n_2}}}\right)^{\alpha_2}$$

$$+M\left|1-\frac{[n_1+1]_{p_{n_1},q_{n_1}}}{c_{n_1}+\gamma_1}\right|^{\alpha_1}\left|1-\frac{[n_2+1]_{p_{n_2},q_{n_2}}}{c_{n_2}+\gamma_2}\right|^{\alpha_2}\left(\frac{p_{n_1}[n_1]_{p_{n_1},q_{n_1}}}{[n_1+1]_{p_{n_1},q_{n_1}}}\frac{x}{1+x}\right)^{\alpha_1}\left(\frac{p_{n_2}[n_2]_{p_{n_2},q_{n_2}}}{[n_2+1]_{p_{n_2},q_{n_2}}}\frac{y}{1+y}\right)^{\alpha_2}+M\delta_{n_1}^{\frac{\alpha_1}{2}}(x)\delta_{n_2}^{\frac{\alpha_2}{2}}(y).$$

This completes the proof. □

References

[1] U. Abel and M. Ivan, Some identities for the operator Bleimann-Butzer and Hahn involving divided differences, Calcolo 36 (3), 1999, 143-160.

[2] U. Abel, On the asymptotic approximation with Bivariate operators of Bleimann-Butzer and Hahn, J. Approx. Theory 97, 1999, 181-198.

[3] U. Abel and M. Ivan, Best constant for a Bleimann-Butzer-Hahn moment estimation, East J. Approx 6 (3), 2000, 1-7.

[4] A. Aral, O. Doğru, Bleimann Butzer and Hahn operators based on q-integers, J. Inequal. Appl., (2007) 1-12. Art. ID 79410.

[5] A. Aral, V. Gupta and R. P. Agarwal. Application of q-Calculus in Operator Theory, Springer, 2013.

[6] G. Bleimann, P.L. Butzer and L. Hahn, A Bernstein-type operator approximating continuous functions on the semi-axis, Indag. Math., 42 (1980) 255-262.

[7] O. Doğru, On Bleimann, Butzer and Hahn type generalization of Balázs operators, Studia Universitatis Babeş-Bolyai. Mathematica, vol. 47, no. 4, pp. 37-45, 2002.

[8] S. Ersan and O. Doğru, Statistical approximation properties of q-Bleimann, Butzer and Hahn operators, Math. Comput. Modell., 49 (2009) 1595-1606.

[9] S. Ersan, Approximation properties of bivariate generalization of Bleimann, Butzer and Hahn operators based on q-integers, 12th WSEAS Int. Conf. on Applied mathematics, Cairo, Egypt, december 29-31, 2007.

[10] S. Ersan and O. Doğru, Statistical approximation properties of q-Bleimann, Butzer and Hahn operators, Math. Comput. Modell., 49 (2009) 1595-1606.

[11] A.D. Gadjiev and Ö. Cakar, On uniform approximation by Bleimann, Butzer and Hahn operators on all positive semi-axis, Trans. Acad. Sci. Azerb.Ser. Phys. Tech. Math. Sci., 19 (1999) 21-26.

[12] B. Lenze, Bernstein-Baskakov-Kantorovich operators and Lipschitz-type maximal functions, in: Colloq. Math. Soc. Janos Bolyai, 58, Approx. Th., (1990) 469-496.

[13] A. Lupaş, A q-analogue of the Bernstein operator, University of Cluj- Napoca, Seminar on Numerical and Statistical Calculus, 9(1987) 85-92.

[14] N.I. Mahmudov, Statistical approximation of Baskakov and Baskakov-Kantorovich operators based on the q-integers, Cent. Eur. Jour. Math., 8(4) (2010) 816-826.

[15] N.I. Mahmudov and V. Gupta, On certain q-analogue of Szász Kantorovich operators, Jour. Appl. Math. Comput., 37 (2011) 407-419.

[16] M. Mursaleen, K. J. Ansari and A. Khan, On (p,q)-analogue of Bernstein Operators, Appl. Math. Comput., 10.1016/j.amc.2015.04.090.

[17] M. Mursaleen, K. J. Ansari and A. Khan,, Some Approximation Results by (p,q)-analogue of Bernstein-Stancu operators, Applied Mathematics and Computation, 10.1016/j.amc.2015.03.135.

[18] M. Mursaleen and A. Khan, Statistical approximation properties of modified q-Stancu-Beta operators, Bull. Malays. Math. Sci. Soc. (2), 36(3) (2013), 683-690.

[19] M. Mursaleen and A. Khan, Generalized q-Bernstein-Schurer operators and some approximation theorems, Jour. Function Spaces Appl., Volume 2013, Article ID 719834, 7 pages.

[20] M. Mursaleen, Md. Nasiruzzaman, Asif Khan and Khursheed J. Ansari, Some approximation results on Bleimann-Butzer-Hahn operators defined by (p,q)-integers, arXiv:1505.00392v1 [math.CA] 3 May 2015.

[21] G.M. Phillips, Bernstein polynomials based on the q-integers, The heritage of P.L.Chebyshev: Ann. Numer. Math. 4 (1997) 511-518.

[22] P. N. Sadjang, On the fundamental theorem of (p,q)-calculus and some (p,q)-Taylor formulas, arXiv:1309.3934 [math.QA].

[23] V. Sahai and S. Yadav, Representations of two parameter quantum algebras and p,q-special functions, J. Math. Anal. Appl. 335 (2007) 268-279.

[24] D. D. Stancu, Approximation of functions by a new class of linear polynomial operators, Revue Roumaine de Mathématiques Pures et Appliquées, vol. 13, pp. 1173-1194, 1968.

[25] D.D. Stancu, A new class of uniform approximating polynomial operators in two and several variables, Proc. Conf. on Constructive Theory of Functions, Budapest (1969), 443-455.

[26] K. Victor and C. Pokman, Quantum Calculus, Springer-Verlag, New York Berlin Heidelberg, 2002.